

## INFLIXIMAB PRESCRIPTION REFERRAL FORM

Memorial Hermann Home Health Pharmacy 21501 Park Row Drive, Suite 210, Katy, Texas 77449 **P** 281.698.6175 **F** 281.698.6147

PATIENT INFORMATI	ON	*Please	include copy of	prescr	iption a	nd medica	linsurance	card, front a	and back*		
Name:								DOB:		☐ Male ☐ Fen	nale
Address:	dress:				City:						
Primary Phone:	rimary Phone:				Alt. Phor	ne:				_	
Email Address:				F	rimary L	_anguage:	☐ English	$\square$ Spanish	☐ Other: _		
PRESCRIBER INFORM	ATION										
Name:					NF	PI:			DEA:		
Name:				City:					State:	Zip:	
Phone:			Fax:				Office C	Contact:			
<b>CLINICAL INFORMAT</b>	ION										
Diagnosis ☐ K50.1 - (ICD-10): ☐ K50.8 - ☐ K50.9 -	Crohn's Disease Crohn's Disease Crohn's Disease	e of the e of Both e, Unspe	h Intestines Coified	☐ M45 ☐ M06 ☐ L40.	.9 - Ank .9 - Rhe 52 - Pso	cylosing Speumatoid A priatic Arth	oondylitis, arthritis, U aritis	Unspecified nspecified	□ L40.9 - F □ Other	Psoriasis Vulgaris Psoriasis, Unspec	ified
Weight: kg He	ight: cm	Date of	Negative TB Tes	st:		Date of Ch	est X-Ray	:  1\	/ Access: $\square$	PIV   Other:	
☐ NKDA ☐ Allergies	s:										
PREVIOUS AND/OR C	URRENT MEDIC	CATIONS	S USED TO TREA	T THI	S DIAGI	NOSIS					
Medication I	Current Start Date			Ind Date Di			iscontinue Reason (if stopped)				
					☐ Failed	☐ Other:					
						☐ Failed ☐ Other:					
					☐ Failed	☐ Other:					
PRESCRIPTION AND	ORDERS										
	☐ No infliximab product preference ☐ Preferred Product:								t dose:		
<b>Infliximab</b> (Remicade, Inflectra, Renflexis, Avsola)	Dosing Regimen Dose				Frequer	псу	су		Quantity/Refills		
	Induction dose	n dose ☐ 3 mg/kg IV			☐ Weeks 0, 2, and 6		☐ 3 doses (infusions)				
	Maintananaa	dooo	☐ 5 mg/kg IV	ŀ						$\dashv$	
	Maintenance (	uose	□ mg/kg IV		☐ Every 8 weeks ☐ Every weeks		☐ doses (infusions) ☐ Fill until follow-up date:				
	Infusion directions: **Do not infuse any other medications along with infliximab**  ☐ Start infusion at 10 mL/hr and increase if tolerated after 15 minutes. Continue to titrate the infusion as tolerated using the following infusion rates: 20 mL/hr x 15 minutes, 40 mL/hr x 15 minutes, 80 mL/hr x 15 minutes, 150 mL/hr x 30 minutes. Maximum infusion rate of 250 mL/hr. Infusion time not less than 2 hours.  ☐ Other:										
Pre-Medications To be administered 30 minutes prior to starting the infusion	□ Acetaminophen: □ 325 mg PO □ 500 mg PO □ 650 mg PO □ Other:mg PO □ Diphenhydramine: □ 25 mg PO □ 50 mg PO □ 25 mg IV □ Other:mg □ PO / □ IV □ Methylprednisolone: □ 40 mg IV □ 125 mg IV □ Other:mg IV □ Other:mg IV										
Adverse Reaction Orders	hypersens hypotensid • Administer	is approjitivity recon, back reaction Diphe   Aceta   One   Norm   Epine	priate and notify pactions to include pain, sudden che management menhydramine 25 maminophen 500 mal Saline 0.9% 50 phrine (1:1,000 snue after receivin	e feve est pa nedica ng IV ng PO 00 mL streng	r, chills, in or hy tions. Othe Othe at a rat th) 0.3 r	dyspnea, pertension er:er:er:er:er g subcut	pruritus, u mg IV PR mg PO PF mL/hr	urticaria, conv N for urticaria RN for myalgia	vulsions, ery a, pruritus, c a or fever gr	thematous rash, r shortness of br eater than 101.3	
Lab Orders	☐ Albumin ☐ Other:	ALT	☐ AST ☐ Creat	tinine	□ CMI			☐ LFT ☐ ☐ Every Infu		er:	
	Nursing to establish and/or maintain venous access, administer prescribed medication, and assess general status and response to therapy.  IV access to be flushed by nurse:  Sodium Chloride 0.9% - 10 mL pre-infusion and 10 mL post-infusion  Other:										
Nursing Orders	☐ Sodiu			L pre-	infusion	and 10 m	L post-infu	ısion		_	
Nursing Orders  Pharmacy Orders	□ Sodiu □ Other	:							omplete the	_ rapy as prescribe	nd.